arXiv:hep-lat/0409077v1 14 Sep 2004

The lattice QCD simulation of the quark-gluon mixed condensate $g\langle\bar{q}\sigma_{\mu\nu}G_{\mu\nu}q\rangle$ at finite temperature and the phase transition of QCD

T. Doi[a]*, N. Ishii[b], M. Oka[b] and H. Suganuma[b]

[a]RIKEN BNL Research Center, Brookhaven National Laboratory, Upton, New York 11973, USA

[b] Dept. of Physics, Tokyo Institute of Technology, Ohokayama 2-12-1, Meguro, Tokyo 152-8551, Japan

The thermal effects on the quark-gluon mixed condensate $g\langle\bar{q}\sigma_{\mu\nu}G_{\mu\nu}q\rangle$, which is another chiral order parameter, are studied using the SU(3)$_c$ lattice QCD with the Kogut-Susskind fermion at the quenched level. We perform the accurate measurement of $g\langle\bar{q}\sigma_{\mu\nu}G_{\mu\nu}q\rangle$ as well as $\langle\bar{q}q\rangle$ for $0 \lesssim T \lesssim 500$MeV. We observe the sharp decrease of both the condensates around $T_c \simeq 280$MeV, while the thermal effects below T_c are found to be weak. We also find that the ratio $m_0^2 \equiv g\langle\bar{q}\sigma_{\mu\nu}G_{\mu\nu}q\rangle/\langle\bar{q}q\rangle$ is almost independent of the temperature even in the very vicinity of T_c, which indicates that the two condensates have nontrivial similarity in the chiral behaviors. We also present the correlation between the condensates and the Polyakov loop to understand the vacuum structure of QCD.

In order to understand the nature of QCD and the relation to hadron phenomenology, the nonperturbative nature is one of the most important characteristics of QCD. The nonperturbative phenomena, such as spontaneous chiral-symmetry breaking and color confinement, also bring the rich structure to the QCD phase diagram. For example, at high temperature, QCD is believed to exhibit phase transition into QGP, where chiral symmetry is restored and the color is deconfined. To realize these phenomena, the RHIC experiments are in progress, which attempt to produce QGP in the laboratory. For the theoretical study of the finite temperature QCD and the phase transition of QCD, we focus on the thermal effects on the condensates. In fact, condensates directly characterize the nontrivial QCD vacuum and thus can indicate the change of the vacuum structure at finite temperature.

Among various condensates, we study the quark-gluon mixed condensate $g\langle\bar{q}\sigma_{\mu\nu}G_{\mu\nu}q\rangle$ at finite temperature as a relevant physical quantity on the chiral structure of the QCD vacuum. Here, the mixed condensate is another chiral order parameter of the QCD vacuum, since the chirality of the quark in $g\langle\bar{q}\sigma_{\mu\nu}G_{\mu\nu}q\rangle$ flips as $g\langle\bar{q}\sigma_{\mu\nu}G_{\mu\nu}q\rangle = g\langle\bar{q}_R(\sigma_{\mu\nu}G_{\mu\nu})q_L\rangle + g\langle\bar{q}_L(\sigma_{\mu\nu}G_{\mu\nu})q_R\rangle$. We emphasize that $g\langle\bar{q}\sigma_{\mu\nu}G_{\mu\nu}q\rangle$ characterizes different aspect of the QCD vacuum from $\langle\bar{q}q\rangle$, because $g\langle\bar{q}\sigma_{\mu\nu}G_{\mu\nu}q\rangle$ reflects the direct correlation between color-octet components of q-$\bar{q}$ pairs and the spontaneously generated gluon field, while $\langle\bar{q}q\rangle$ reflects only the color-singlet q-$\bar{q}$ components.

We further note that the mixed condensate is important quantity in hadron phenomenology through the QCD sum rule framework. In fact, it is known that $g\langle\bar{q}\sigma_{\mu\nu}G_{\mu\nu}q\rangle$ has large effects in the QCD sum rule especially for baryons, such as N-Δ splitting [1] and parity splitting [2]. Recently, it is also shown [3] that $g\langle\bar{s}\sigma_{\mu\nu}G_{\mu\nu}s\rangle/\langle\bar{s}s\rangle$ is a key quantity for the prediction on the parity of the penta-quark baryon, $\Theta^+(1540)$ [4].

To study the thermal effects on $g\langle\bar{q}\sigma_{\mu\nu}G_{\mu\nu}q\rangle$, we use lattice QCD Monte Carlo simulation, which is the direct and nonperturbative calculation from QCD. So far, the lattice QCD studies for $g\langle\bar{q}\sigma_{\mu\nu}G_{\mu\nu}q\rangle$ have been performed at zero temperature by a pioneering but rather preliminary work [5], and by the recent works of our group [6] using the KS fermion, and of another group [7] using the Domain-Wall fermion. At finite temperature, however, there has been no result on $g\langle\bar{q}\sigma_{\mu\nu}G_{\mu\nu}q\rangle$ except for our early reports [8]. Therefore, we present the extensive results of the thermal effects on $g\langle\bar{q}\sigma_{\mu\nu}G_{\mu\nu}q\rangle$ with the analysis near the critical temperature [9].

We evaluate the condensates $g\langle\bar{q}\sigma_{\mu\nu}G_{\mu\nu}q\rangle$ as well as $\langle\bar{q}q\rangle$ using the $\mathrm{SU}(3)_c$ lattice QCD at the quenched level. The Monte Carlo simulations are performed with the standard Wilson action for $\beta = 6.0, 6.1$ and 6.2. The lattice units are obtained as $a^{-1} = 1.9$, 2.3 and 2.7GeV for $\beta =$ 6.0, 6.1 and 6.2, respectively, so as to reproduce the string tension $\sqrt{\sigma} = 427$MeV [9]. We perform the calculation at various temperatures as $0 \lesssim T \lesssim 500$MeV using the following lattices,

---

*e-mail: doi@quark.phy.bnl.gov; The Monte Carlo simulations have been performed on NEC SX-5 at Osaka University and IBM Regatta at Tokyo Institute of Technology.

(i) $\beta = 6.0$, $16^3 \times N_t$ $(N_t = 16, 12, 10, 8, 6, 4)$,
(ii) $\beta = 6.1$, $20^3 \times N_t$ $(N_t = 20, 12, 10, 8, 6)$,
(iii) $\beta = 6.2$, $24^3 \times N_t$ $(N_t = 24, 16, 12, 10, 8)$.

We generate 100 gauge configurations for each lattice. Moreover, we generate 1000 gauge configurations in the vicinity of the phase transition point, namely, $20^3 \times 8$ at $\beta = 6.1$ and $24^3 \times 10$ at $\beta = 6.2$, because the fluctuations of the condensates get larger near T_c. For the lattices at $T > T_c$, we only use the gauge configurations which are continuously connected to the trivial vacuum $U_\mu = 1$. For the fermion action, we employ the KS fermion to preserve the explicit chiral symmetry for the quark mass $m = 0$, which is desirable for the study of chiral order parameters.

In the calculation of the condensates on the lattice, we use the formula as $a^3\langle\bar{q}q\rangle = -\frac{1}{4}\sum_f \mathrm{Tr}\left[\langle q^f(x)\bar{q}^f(x)\rangle\right]$, $a^5 g\langle\bar{q}\sigma_{\mu\nu}G_{\mu\nu}q\rangle = -\frac{1}{4}\sum_{f,\,\mu,\nu} \mathrm{Tr}\left[\langle q^f(x)\bar{q}^f(x)\rangle\sigma_{\mu\nu}G^{\mathrm{lat}}_{\mu\nu}(x)\right]$, where $\mathrm{SU}(4)_f$ quark-spinor fields, q and $\bar{q}$, are converted into spinless Grassmann KS fields and the gauge-link variable in the actual calculations. The more detailed formula are given in Ref. [6].

In each configuration, we measure the condensates on 16 ($\beta = 6.0$) or 2 ($\beta = 6.1, 6.2$) different points which are taken so as to be equally spaced on the 4-dimensional lattice [6,9]. Therefore, we achieve high statistics as 1600 data at $\beta = 6.0$ and 200 data at $\beta = 6.1$ and 6.2. Note that for the lattices of $20^3 \times 8$ ($\beta = 6.1$) and $24^3 \times 10$ ($\beta = 6.2$), we obtain 2000 data, which guarantees reliability of the results even in the vicinity of T_c.

We calculate the condensates at each quark mass of $m \simeq 20, 35, 50$ MeV. At each temperature, we observe that both the condensates show a clear linear behavior against m, and therefore we fit the data with a linear function and determine the condensates in the chiral limit. We estimate the statistical error to be 5-7% level using jackknife error method. The finite-volume artifact is estimated to be about 1% level, through a check of boundary condition effects [6,9].



Figure 1. $g\langle\bar{q}\sigma_{\mu\nu}G_{\mu\nu}q\rangle_T$ normalized by $g\langle\bar{q}\sigma_{\mu\nu}G_{\mu\nu}q\rangle_{T=0}$ plotted against temperature T. The vertical dashed line denotes the critical temperature



Figure 2. $m_0^2(T) \equiv g\langle\bar{q}\sigma_{\mu\nu}G_{\mu\nu}q\rangle_T / \langle\bar{q}q\rangle_T$ normalized by $m_0^2(T = 0)$ plotted against T. This result indicates the same chiral behavior between $g\langle\bar{q}\sigma_{\mu\nu}G_{\mu\nu}q\rangle_T$ and $\langle\bar{q}q\rangle_T$.

We evaluate the thermal effects on each condensate, $g\langle\bar{q}\sigma_{\mu\nu}G_{\mu\nu}q\rangle$ or $\langle\bar{q}q\rangle$, by taking the ratio between the values at finite and zero temperatures. Note that the renormalization constants cancel in these ratios. In figure 1, we plot the thermal effects on $g\langle\bar{q}\sigma_{\mu\nu}G_{\mu\nu}q\rangle$. We find a drastic change of $g\langle\bar{q}\sigma_{\mu\nu}G_{\mu\nu}q\rangle$ around the critical temperature $T_c \simeq 280$MeV. We obtain $T_c/\sqrt{\sigma} = 0.64(4)$, which is consistent with the coincidence of the confinement/deconfinement phase transition and chiral-symmetry restoration. We also find that the thermal effects on $g\langle\bar{q}\sigma_{\mu\nu}G_{\mu\nu}q\rangle$ are remarkably weak for $T \lesssim 0.9\,T_c$. The same nontrivial features are also found for $\langle\bar{q}q\rangle$.

We then quantitatively compare the thermal effects of $g\langle\bar{q}\sigma_{\mu\nu}G_{\mu\nu}q\rangle$ and $\langle\bar{q}q\rangle$ by plotting the ratio $m_0^2(T) \equiv g\langle\bar{q}\sigma_{\mu\nu}G_{\mu\nu}q\rangle_T/\langle\bar{q}q\rangle_T$ against T. In figure 2, we observe that $m_0^2(T)$ is almost in-



Figure 3. Time history of $g\langle\bar{q}\sigma_{\mu\nu}G_{\mu\nu}q\rangle_T$ (upper) and the Polyakov loop (lower) for the 500 configurations on $24^3 \times 10$ ($\beta = 6.2$) lattice near T_c. This result indicates strong correlation between these two quantities.

dependent of the temperature, even in the very vicinity of T_c. This nontrivial result can be interpreted that $g\langle\bar{q}\sigma_{\mu\nu}G_{\mu\nu}q\rangle_T$ and $\langle\bar{q}q\rangle_T$ have similarity in the chiral behaviors.

To understand this nontrivial similarity, we express the condensates as $\langle\bar{q}q\rangle = \frac{1}{V}\int d\lambda' \frac{m\rho(\lambda')}{\lambda'^2+m^2}$, $g\langle\bar{q}\sigma_{\mu\nu}G_{\mu\nu}q\rangle = \frac{1}{V}\int d\lambda' \frac{m\rho(\lambda')}{\lambda'^2+m^2}\langle\lambda'|\sigma_{\mu\nu}G_{\mu\nu}|\lambda'\rangle$, where $|\lambda\rangle$ denotes the eigenvector of the Dirac operator and $\rho(\lambda)$ the spectral density on λ [10]. Using these formula, the common thermal behavior is understood in the way that the thermal effects are dominated by $\rho(\lambda)|_{\lambda=0}$ and $\langle\lambda|\sigma_{\mu\nu}G_{\mu\nu}|\lambda\rangle|_{\lambda=0}$ has remarkably weak dependence on T.

This result further implies that the thermal effects hardly appear in the local gluon field strength, but appear in the global structure of the vacuum such as topological quantities. To proceed, we plot the time history of $g\langle\bar{q}\sigma_{\mu\nu}G_{\mu\nu}q\rangle$ and the Polyakov loop in figure 3, and find the strong correlation between $g\langle\bar{q}\sigma_{\mu\nu}G_{\mu\nu}q\rangle$ and the Polyakov loop. For the gauge configuration with a small (large) value of $g\langle\bar{q}\sigma_{\mu\nu}G_{\mu\nu}q\rangle$, the Polyakov loop takes a large (small) value. One of the possible explanation is that the thermal effects are dominated by the quark propagation making a circuit in time direction, which can directly reflect the topological structure of the vacuum.

In summary, we have studied thermal effects on $g\langle\bar{q}\sigma_{\mu\nu}G_{\mu\nu}q\rangle$ using the SU$(3)_c$ lattice QCD with the KS fermion at the quenched level. We have observed a clear signal of chiral restoration as a sharp decrease of $g\langle\bar{q}\sigma_{\mu\nu}G_{\mu\nu}q\rangle$ as well as $\langle\bar{q}q\rangle$, while the thermal effects have been found to be small for $T \lesssim 0.9\,T_c$. We have also found that $m_0^2(T) \equiv g\langle\bar{q}\sigma_{\mu\nu}G_{\mu\nu}q\rangle_T/\langle\bar{q}q\rangle_T$ is almost independent of T in the entire region up to T_c, which indicates that these chiral condensates show a common thermal behavior. The strong correlation between the condensates and the Polyakov loop has been also observed. For further studies, full QCD lattice calculations are interesting to analyze dynamical quark effects on the condensates.

REFERENCES

1. H.G. Dosch, M. Jamin, and S. Narison, Phys. Lett. **B220** (1989) 251; W-Y.P. Hwang and K.-C. Yang, Phys. Rev. **D49** (1994) 460.
2. D. Jido, N. Kodama and M. Oka, Phys. Rev. **D54** (1996) 4532; D. Jido and M. Oka, hep-ph/9611322.
3. J. Sugiyama, T. Doi and M. Oka, Phys. Lett. **B581** (2004) 167.
4. T. Nakano et al., Phys. Rev. Lett. **91** (2003) 012002.
5. M. Kremer and G. Schierholz, Phys. Lett. **B194** (1987) 283 .
6. T. Doi, N. Ishii, M. Oka and H. Suganuma, Phys. Rev. **D67** (2003) 054504; Proc. of "Quark Confinement and the Hadron Spectrum V", (World Scientific, 2003) p.381
7. T.W. Chiu and T.H. Hsieh, Nucl. Phys. **B673** (2003) 217.
8. T. Doi, N. Ishii, M. Oka, H. Suganuma, Nucl. Phys. **A721** (2003) 934; Prog. Theor. Phys. Suppl. **151** (2003) 161; Nucl. Phys. **B129** (Proc. Suppl.) (2004) 566; Proc. of "Color Confinement and Hadrons in Quantum Chromodynamics", (World Scientific, 2004) p.398.
9. T. Doi, N. Ishii, M. Oka and H. Suganuma, Phys. Rev. **D70** (2004) 034510.
10. T. Banks and A. Casher, Nucl. Phys. **B169** (1980) 103; S. Hands, J.B. Kogut and A. Kocic, Nucl. Phys. **B357** (1991) 467.